

<i>recursion scheme</i>	<i>adjunction</i>	<i>conjugates</i>	<i>para-hylo equation</i>	<i>algebra</i>
(hylo-shift law)	$\text{Id} \dashv \text{Id}$	$\alpha \dashv \alpha$	$x = a \cdot (\text{id} \triangle \mathsf{D} x \cdot \alpha C \cdot c) : A \leftarrow C$	$a : C \times \mathsf{D} A \rightarrow A$
mutual recursion	$\Delta \dashv (\times)$	ccf	$x_1 = a_1 \cdot (\text{id} \triangle \mathsf{D} (x_1 \triangle x_2) \cdot c) : A_1 \leftarrow C$ $x_2 = a_2 \cdot (\text{id} \triangle \mathsf{D} (x_1 \triangle x_2) \cdot c) : A_2 \leftarrow C$	$a_1 : C \times \mathsf{D} (A_1 \times A_2) \rightarrow A_1$ $a_2 : C \times \mathsf{D} (A_1 \times A_2) \rightarrow A_2$
accumulator	$- \times P \dashv (-)^P$	ccf	$x = a \cdot (\text{outl} \triangle ((\mathsf{D} (\wedge x) \cdot c) \times P)) : A \leftarrow C \times P$	$a : C \times \mathsf{D} (A^P) \times P \rightarrow A$
course-of-values (§5.6)	$\mathsf{U}_{\mathsf{D}} \dashv \mathsf{Cofree}_{\mathsf{D}}$	ccf	$x = a \cdot (\text{id} \triangle \mathsf{D} (\mathsf{D}_{\infty} x \cdot \mathbf{[c]}) \cdot c) : A \leftarrow C$	$a : C \times \mathsf{D} (\mathsf{D}_{\infty} A) \rightarrow A$
finite memo-table (§5.6)	$\mathsf{U}_{*} \dashv \mathsf{Cofree}_{*}$	ccf	$x = a \cdot (\text{id} \triangle \mathsf{D} (\mathsf{D}_{*} x \cdot \mathbf{[c]_{*}}) \cdot c) : A \leftarrow C$	$a : C \times \mathsf{D} (\mathsf{D}_{*} A) \rightarrow A$

Table 1. Different types of para-hylos building on the canonical control functor (ccf); the coalgebra is $c : C \rightarrow \mathsf{D} C$ in each case.